arXiv:1602.04093v1 [math.GR] 12 Feb 2016

Enumerating fibres of commutator words over p-groups

Matthew Levy

Bielefeld University

Abstract

We enumerate the fibres of commutator word maps over p-groups of nilpotency class less than p with exponent p. We also give some examples and enumerate the fibre sizes of all word maps over p-groups of class 2 with exponent p.

1 Introduction

Let G be a finite group, $w(x_1, ..., x_n)$ a group word and w the associated word map $w : G^{(n)} \to G$. For $g \in G$ denote by $N_w^G(g)$ the number of solutions to $w = g$ and by $P_w^G(g)$ the probability that a random n-tuple $\mathbf{g} = (g_1, ..., g_n) \in G^{(n)}$ satisfies $w(\mathbf{g}) = g$, i.e.

$$P_w^G(g) = \frac{N_w^G(g)}{|G|^n}.$$

The study of P_w^G has attracted a lot of attention in recent years. For example, in [4], it was shown that for $1 \neq w$ and a finite simple group G, $P_w^G(1) \to 1$ as $|G| \to \infty$. Results in [10] provide sharp bounds on P_w^G for general words w and finite simple groups G. If G is abelian, then the word map

$$w : G^{(n)} \to G,$$

is a homomorphism and it is clear that

$$N_w^G(1) = |\text{Ker } w| = \frac{|G|^n}{|\text{Im } w|} \geq |G|^{n-1}$$

and so $P_w^G(1) \geq \frac{1}{|G|}$. It is a conjecture of Alon Amit (see [1]) that if G is a nilpotent group then $P_w^G(1) \geq \frac{1}{|G|}$. In [11] we prove Amit's Conjecture in the special case where the nilpotency class is 2. Note that since the statistics $N_w^G(1)$ and $P_w^G(1)$ are multiplicative under direct products, we may reduce to the case where G is a finite p-group. In this paper we study the fibre sizes of a particular class of words over p-groups. In Section 3 we show that this is enough to determine the fibre sizes of all word maps over p-groups of class 2 with exponent p.

For $t \in \mathbb{N}$, let c_t denote the word map given by $c_t = [x_1, y_1]...[x_t, y_t]$ and, for $g \in G$, let $N_t^G(g)$ denote $N_{c_t}^G(g)$. Similarly let P_t^G denote the *c_t-distribution* on

G, i.e.

$$P_t^G(g) = \frac{N_t^G(g)}{|G|^{2t}},$$

and let U^G be the uniform distribution on G (i.e. $U^G(g) = 1/|G|$). By a classical result of Frobenius from 1896 (see, for example, [7]) we have

$$N_t^G(g) = |G|^{2t-1} \sum_{\chi \in \mathrm{Irr}(G)} \frac{\chi(g)}{\chi(1)^t}, \tag{1}$$

where 1 is the identity element of G and we sum over the irreducible complex characters of G. In the main result of this paper, Theorem 2.1, we give explicit formulae to compute the numbers $N_t^G(g)$, for finite p-groups G of nilpotency class less than p with exponent p, in terms of the number of rational points of certain algebraic varieties.

In [5] Garion & Shalev prove the following.

Proposition 1.1 (Proposition 1.1 [5])**.** *Let G be a finite group. Then*

$$||P_1^G - U^G||_1 \leq \left(\sum_{\chi \in \mathrm{Irr}(G), \chi \neq Id} \chi(1)^{-2} \right)^{1/2},$$

where $||P_1^G - U^G||_1 = \sum_{g \in G} |P_1^G(g) - U^G(g)|$.

As the authors remark, this bound has no content when the sum on the right hand side is greater than or equal to 1. Since the non-trivial linear characters of G contribute $|G/G'|^{-1}$ to the sum the result can only be useful for perfect groups. Since the maps c_t take values in G' it is not hard to adapt their proof and deduce the following.

Proposition 1.2. *Let G be a finite group. Then*

$$||P_t^{G'} - U^{G'}||_1 \leq \left(\frac{|G'|}{|G|} \sum_{\chi \in \mathrm{Irr}(G), \chi(1) \neq 1} \chi(1)^{-2t} \right)^{1/2}.$$

If the right hand side in Proposition 1.2 is close to zero, then the fibres of the maps c_t are roughly the same size and the values are uniformly distributed. The proof follows easily from the following lemmas.

Lemma 1.3. *Let G be a finite group. Then*

$$\sum_{g \in G'} P_t^G(g)^2 = \frac{1}{|G'|} + \frac{1}{|G|} \sum_{\chi \in \mathrm{Irr}(G), \chi(1) \neq 1} \chi(1)^{-2t}.$$

Proof. The proof follows from Lemma 2.1 in [5] and noting that $P_t^G(g) = 0$ for $g \notin G'$. □

Lemma 1.4. *Let G be a finite group. Then*

$$\sum_{g \in G'} \left(P_t^G(g) - \frac{1}{|G'|} \right)^2 = \frac{1}{|G|} \sum_{\chi \in \mathrm{Irr}(G), \chi(1) \neq 1} \chi(1)^{-2t}.$$

Proof. By Lemma 1.3,

$$\begin{aligned}\sum_{g\in G'}\left(P_t^G(g)-\frac{1}{|G'|}\right)^2 &= \sum_{g\in G'}P_t^G(g)^2-\frac{2}{|G'|}\sum_{g\in G'}P_t^G(g)+\frac{1}{|G'|}\\ &= \frac{1}{|G|}\sum_{\chi\in\mathrm{Irr}(G),\chi(1)\neq 1}\chi(1)^{-2t}\end{aligned}$$

since $\sum_{g\in G'}P_t^G(g)=1$. □

Proof of Proposition 1.2. This follows from the Cauchy-Schwarz inequality,

$$(||P_t^{G'}-U^{G'}||_1)^2=\left(\sum_{g\in G'}(P_t(g)-\frac{1}{|G'|}\right)^2\leq |G'|\sum_{g\in G'}\left(P_t(g)-\frac{1}{|G'|}\right)^2$$

and the previous lemma. □

In Section 2 we will develop formulae for the fibre sizes of the word maps c_t over p-groups of nilpotency class less than p with exponent p. Moreover, these results extend, more generally, to p-groups obtained by 'base extension'. In Section 3 we will determine the fibre sizes of all word maps over p-groups of class 2 with exponent p. We will then give some examples in Section 4.

2 Enumerating sizes of fibres

Let G be a finite p-group and, for each $i\in\mathbb{N}$, write $\mathrm{Irr}^i(G)=\{$irreducible complex characters of G of degree $p^i\}$ and $\mathrm{Irr}(G)$ for the set of all irreducible complex characters. For a complex variable s and $g\in G$ write

$$\zeta_G^i(s,g)=\sum_{\chi\in\mathrm{Irr}^i(G)}\frac{\chi(g)}{\chi(1)^s}. \tag{2}$$

We will also write

$$\zeta_G(s,g)=\sum_{\chi\in\mathrm{Irr}(G)}\frac{\chi(g)}{\chi(1)^s}=\sum_{i\in\mathbb{N}}\zeta_G^i(s,g). \tag{3}$$

It is clear from equations (1) and (3) that

$$N_t^G(g)=|G|^{2t-1}\zeta_G(t,g)$$

and that

$$P_t^G(g)=\frac{1}{|G|}\zeta_G(t,g). \tag{4}$$

In this section we prove Theorem 2.1 which allows us to compute expressions like equations (3) and (4) for finite p-groups of nilpotency class less than p with exponent p.

Note that when $s=1$ and $g=1$, the identity of G, the sum $\zeta_G(1,1)$ is simply the class number $k(G)$ of G and that if g is not in the derived group of G then $\zeta_G(s,g)=0$ since $N_s^G(g)=0$, see equation (1). An analogue of equation

(3), a *twisted* zeta function, is studied by Jaikin-Zapirain c.f. [8, Theorem 1.2] where he shows that it is a rational function in p^{-s} for any $g \in G$ where G is a FAb uniform pro-p group.

We also note that the twisted zeta function in equation (3) is multiplicative under direct products in the following sense: for $g_1 \in G_1$ and $g_2 \in G_2$ where G_1 and G_2 are groups we have

$$\zeta_{G_1 \times G_2}(s, g_1 g_2) = \zeta_{G_1}(s, g_1).\zeta_{G_2}(s, g_2).$$

In [12] O'Brien & Voll use the Kirillov orbit method to enumerate the irreducible complex characters, of each degree, of finite p-groups of nilpotency class less than p. If the group G is of exponent p, then the number of characters, of each degree, of G can be described in terms of the number of rational points of certain algebraic varieties. We can analogously compute the sums $\zeta_G^i(s, g)$ by counting the numbers of rational points of the algebraic varieties considered by O'Brien & Voll that intersect a hyperplane characterized by g.

We now fix a p-group G of nilpotency class less than p with exponent p and an element $g \in G'$.

The Lazard correspondence establishes an order-preserving bijection between finite p-groups of nilpotency class $c < p$ and finite nilpotent Lie rings of p-power order and class $c < p$; cf. [9, Example 10.24]. Let $c < p$ be the nilpotency class of G. Let $\mathfrak{g} = \log(G)$ be the finite Lie ring associated to G by the Lazard correspondence. The Kirillov orbit method gives a correspondence between characters of G and orbits in $\hat{\mathfrak{g}} := \mathrm{Hom}_{\mathbb{Z}}(\mathfrak{g}, \mathbb{C}^*)$, the Pontryagin dual of $\mathfrak{g}$, under the co-adjoint action of G on $\hat{\mathfrak{g}}$; cf. [3, Theorem 2.6] or [6, Theorem 4.4]. Under this correspondence each orbit Ω of size, say, p^{2i} gives rise to a character χ_Ω of degree p^i and all characters are of this form. We have

$$\zeta_G^i(s, g) = \sum_{\Omega \subseteq \hat{\mathfrak{g}}, |\Omega| = p^{2i}} \frac{\chi_\Omega(g)}{\chi_\Omega(1)^s},$$

where we sum over orbits Ω of $\hat{\mathfrak{g}}$ and χ_Ω is the character of G that corresponds to the orbit Ω. For each $\omega \in \Omega$ we denote $\chi_\omega = \chi_\Omega$.

For each $\omega \in \hat{\mathfrak{g}}$ we write B_ω for the bi-additive, skew-symmetric form $B_\omega : \mathfrak{g} \times \mathfrak{g} \to \mathbb{C}^*$, $(u, v) \mapsto \omega([u, v])$ and $\mathrm{Rad}(B_\omega)$ for the radical of B_ω. Note that the centre $\mathfrak{z}$ of $\mathfrak{g}$ is contained in $\mathrm{Rad}(B_\omega)$ and the form B_ω only depends on the restriction of ω to $\mathfrak{g}'$. From [12] we have, for each i,

$$\begin{aligned}
\zeta_G^i(s, g) &= p^{-2i} \sum_{\omega \in \hat{\mathfrak{g}}, |\mathfrak{g} : \mathrm{Rad}(B_\omega)| = p^{2i}} \frac{\chi_\omega(g)}{\chi_\omega(1)^s} \\
&= |\mathfrak{g}/\mathfrak{g}'| p^{-2i} \sum_{\omega \in \widehat{\mathfrak{g}'}, |\mathfrak{g} : \mathrm{Rad}(B_\omega)| = p^{2i}} \frac{\chi_\omega(g)}{\chi_\omega(1)^s} \\
&= |\mathfrak{g}/\mathfrak{g}'| p^{-2i} \sum_{\omega \in \widehat{\mathfrak{g}'}, |\,\mathrm{Rad}(B_\omega) : \mathfrak{z}| = p^{-2i} |\mathfrak{g}/\mathfrak{z}|} \frac{\chi_\omega(g)}{\chi_\omega(1)^s}.
\end{aligned}$$

By the Kirillov orbit method we have, for all $\omega \in \hat{\mathfrak{g}}$ such that $|\mathfrak{g} : \mathrm{Rad}(B_\omega)| = p^{2i}$,

$$\chi_\omega(g) = p^{-i} \sum_{v \in \Omega_\omega} v(g),$$

where Ω_ω is the orbit containing ω and we identify g with an element of $\mathfrak{g} = \log(G)$. Hence

$$\zeta_G^i(s,g) = |\mathfrak{g}/\mathfrak{g}'|p^{-i(3+s)} \sum_{\omega \in \widehat{\mathfrak{g}'}, |\operatorname{Rad}(B_\omega):\mathfrak{z}| = p^{-2i}|\mathfrak{g}/\mathfrak{z}|} \sum_{\upsilon \in \Omega_\omega} \upsilon(g).$$

Theorem B in [12] gives a geometric characterization of this sum in terms of the numbers of certain rational points of rank varieties of matrices of linear forms. Assume that $\mathfrak{o}$ is a compact discrete valuation ring of characteristic zero with maximal ideal $\mathfrak{p}$ and residue field $\mathbf{k} = \mathfrak{o}/\mathfrak{p}$ of characteristic p. Suppose that $\mathfrak{g}$ is a finite, nilpotent $\mathfrak{o}$-Lie algebra of class $c < p$ and that both $\mathfrak{g}/\mathfrak{z}$ and $\mathfrak{g}'$ are annihilated by $\mathfrak{p}$. Set $a := \operatorname{rk}_{\mathbf{k}}(\mathfrak{g}/\mathfrak{z})$ and $b := \operatorname{rk}_{\mathbf{k}}(\mathfrak{g}')$ and fix bases $\mathbf{e} = \{e_1, ..., e_a\}$ and $\mathbf{f} = \{f_1, ..., f_b\}$ for $\mathfrak{g}/\mathfrak{z}$ and $\mathfrak{g}'$ respectively. Choose structure constants $\lambda_{ij}^k \in \mathbf{k}$ such that

$$[e_i, e_j] = \sum_{k=1}^{b} \lambda_{ij}^k f_k$$

and $\lambda_{ij}^k = -\lambda_{ji}^k$ for all $i, j \in \{1, ..., a\}$ and $k \in \{1, ..., b\}$. Let $\mathbf{Y} = (Y_1, ..., Y_b)$ be independent variables and define the 'commutator matrix' (with respect to $\mathbf{e}$ and $\mathbf{f}$) $B(\mathbf{Y}) \in \operatorname{Mat}_a(\mathbf{k}[\mathbf{Y}])$ given by

$$B(\mathbf{Y})_{ij} := \sum_{k=1}^{b} \lambda_{ij}^k Y_k$$

for all $i, j \in \{1, ..., a\}$. For $\mathbf{y} = (y_1, ..., y_b) \in \mathbf{K}^b$, where $\mathbf{K}$ is any finite extension of $\mathbf{k}$, write $B(\mathbf{y}) \in \operatorname{Mat}_a(\mathbf{K})$ for the matrix obtained by evaluating the variables Y_i at y_i. Note that the matrices $B(\mathbf{y})$ are skew-symmetric, have even rank and that $\det(B(\mathbf{Y}))$ is a square in $\mathbf{k}[\mathbf{Y}]$, whose square root $\operatorname{Pf}(B(\mathbf{Y})) := \sqrt{\det(B(\mathbf{Y}))}$ is the *Pfaffian* of $B(\mathbf{Y})$. If a is odd then $\operatorname{Pf}(B(\mathbf{Y})) = 0$.

Fix a non-trivial additive character $\phi : \mathbf{k} \to \mathbb{C}^*$. For $a \in \mathbf{k}$ define $\phi_a(x) = \phi(ax)$. The map $a \mapsto \phi_a$ is an isomorphism between $\mathbf{k}$ and its Pontryagin dual $\widehat{\mathbf{k}}$. We have an isomorphism between $\mathfrak{g}'$ and its dual $\widehat{\mathfrak{g}'}$ and also a canonical isomorphism between $\mathfrak{g}'$ and its linear dual $\operatorname{Hom}_{\mathbb{F}_q}(\mathfrak{g}', \mathbf{k})$. We now fix an isomorphsim $\psi_1 : \widehat{\mathfrak{g}'} \to \operatorname{Hom}_{\mathbf{k}}(\mathfrak{g}', \mathbf{k})$. The dual $\mathbf{k}$-basis $f^\vee = (f_k^\vee)$ for $\operatorname{Hom}_{\mathbf{k}}(\mathfrak{g}', \mathbf{k})$ gives a coordinate system

$$\begin{aligned} \psi_2 : \operatorname{Hom}_{\mathbf{k}}(\mathfrak{g}', \mathbf{k}) &\to \mathbf{k}^b; \\ y = \sum_{k=1}^{b} y_k f_f^\vee &\mapsto \mathbf{y} = (y_1, ..., y_b). \end{aligned}$$

Set $\psi := \psi_2 \circ \psi_1 : \widehat{\mathfrak{g}'} \to \mathbf{k}^b$, an isomorphism. Under this isomorphism we may identify elements $\mathbf{y} \in \mathbf{k}^b$ with elements $\omega_{\mathbf{y}} \in \widehat{\mathfrak{g}'}$.

Assume that $\mathfrak{O}$ is an unramified extension of $\mathfrak{o}$, with maximal ideal $\mathfrak{P}$. Write $\mathfrak{g}(\mathfrak{O})$ for $\mathfrak{g} \otimes_{\mathfrak{o}} \mathfrak{O}$ and $\mathfrak{z}(\mathfrak{O})$ for $\mathfrak{z} \otimes_{\mathfrak{o}} \mathfrak{O}$. We identify the residue field $\mathfrak{O}/\mathfrak{P}$, a finite extension of $\mathbf{k}$, with $\mathbb{F}_q$. The derived $\mathfrak{O}$-Lie algebra $\mathfrak{g}(\mathfrak{O})'$ as well as the quotient $\mathfrak{g}(\mathfrak{O})/\mathfrak{z}(\mathfrak{O})$ are annihilated by $\mathfrak{P}$. We denote the corresponding bases, $\mathbf{e} \otimes_{\mathbf{k}} 1$ for $\mathfrak{g}(\mathfrak{O})/\mathfrak{z}(\mathfrak{O})$ and $\mathbf{f} \otimes_{\mathbf{k}} 1$ for $\mathfrak{g}(\mathfrak{O})'$, obtained by this base extension by $\mathbf{e}$ and

f respectively. We consider **e** and **f** as $\mathbb{F}_q$ bases for the respective $\mathbb{F}_q$-vector spaces of dimensions a and b. The group $G = G(\mathfrak{O}) := \exp(\mathfrak{g}(\mathfrak{O}))$ has the property that both $G(\mathfrak{O})'$ and $G(\mathfrak{O})/Z(G(\mathfrak{O}))$ have exponent p. It follows from [12, Theorem B] that

$$\zeta_G^i(s,g) = |\mathfrak{g}/\mathfrak{g}'|p^{-i(3+s)} \sum_{\mathbf{y}\in\mathbb{F}_q^b, \mathrm{rk}(B(\mathbf{y}))=2i} \sum_{\mathbf{v}\in\Omega_{\omega_\mathbf{y}}} \mathbf{v}(g).$$

Since $|\Omega_{\omega_\mathbf{y}}| = p^{2i}$ we have

$$\zeta_G^i(s,g) = |\mathfrak{g}/\mathfrak{g}'|p^{-i(1+s)} \sum_{\mathbf{y}\in\mathbb{F}_q^b, \mathrm{rk}(B(\mathbf{y}))=2i} \omega_\mathbf{y}(g).$$

Let

$$K_G^i(g) = \#\{\mathbf{y}\in\mathbb{F}_q^b : \mathrm{rk}(B(\mathbf{y})) = 2i, g\in\mathrm{Ker}(\omega_\mathbf{y})\} \tag{5}$$

and

$$V_G^i(g) = \#\{\mathbf{y}\in\mathbb{F}_q^b : \mathrm{rk}(B(\mathbf{y})) = 2i, \mathrm{ord}(\omega_\mathbf{y}(g)) \neq 1\}. \tag{6}$$

Note that $\sum_i K_G^i(1) = q^b$ and $V_G^i(1) = 0$ for all i whilst for $1 \neq g \in G$ we have $\sum_i K_G^i(g) = q^{b-1}$.

We write $K_G(g)$ and $V_G(g)$ for the vectors $(K_G^i(g))_i$ and $(V_G^i(g))_i$ respectively. The numbers $\omega_\mathbf{y}(g)$ are p-th roots of unity. Since the sum of the $q-1$ Galois conjugates of non-trivial q-th roots of unity is -1 we have the following:

Theorem 2.1. *Let $\mathfrak{o}$ be a compact discrete valuation ring of characteristic zero with residue field $\mathbf{k}$ of characteristic p and let $\mathfrak{g}$ be a finite, nilpotent $\mathfrak{o}$-Lie algebra of class $c < p$. Assume that $\mathfrak{g}' \cong \mathbf{k}^b$ and $\mathfrak{g}/\mathfrak{z} \cong \mathbf{k}^a$ as $\mathbf{k}$-vector spaces. Let $\mathfrak{O}$ be a finite, unramified extension of $\mathfrak{o}$, with residue field isomorphic to $\mathbb{F}_q$. Write $G := G(\mathfrak{O})$ for the group associated with $\mathfrak{g}(\mathfrak{O})$ under the Lazard correspondence. Then, for each i,*

$$\zeta_G^i(s,g) = |G/G'|p^{-i(1+s)}(K_G^i(g) - \frac{1}{q-1}V_G^i(g)),$$

where K_G^i and V_G^i are defined in equations (5) and (6).

This follows immediately from the discussion above.

3 Nilpotent class 2 groups

Let G be a finite group, $w(x_1,...,x_n)$ a group word and w the associated word map. Recall that $N_w^G(g)$ is the number of solutions to $w = g$ and that $P_w^G(g)$ is the probability that a random n-tuple $\mathbf{g} = (g_1,...,g_n) \in G^{(n)}$ satisfies $w(\mathbf{g}) = g$. The following corollary follows immediately from the results in [11]:

Corollary 3.1 ([11])**.** *Let G be a finite p-group of nilpotency class 2 with exponent p and let w be a group word. Then there exists a word v of the following form:*

i) $v = x$; or

ii) $v = c_t$ for some t,

such that $P_w^G(g) = P_v^G(g)$ *for all* $g \in G$.

By Theorem 2.1 we have determined the fibre sizes of all word maps over all p-groups of nilpotency class 2 with exponent p:

Corollary 3.2. *Let p be an odd prime and let G be a finite p-group of nilpotency class 2 with exponent p and let w be a group word. Then either*

i) $P_w^G(g) = \frac{1}{|G|}$ *for all* $g \in G$*; or*

ii) $P_w^G(g) = \frac{1}{|G|}\zeta_G(t, g)$ *for some t depending on w.*

4 Examples

We compute the sums $\zeta_G^i(s, g)$ for various relatively free p-groups with exponent p. Note that for $i = 0$, the only vector $\mathbf{y}$ giving rise to a matrix $B(\mathbf{y})$ such that $\mathrm{rk}(B(\mathbf{y})) = 0$ (see equations (5) and (6)) is $\mathbf{y} = \mathbf{0}$. Suppose that $i \neq 0$. Since the rank of the matrix $B(\mathbf{y})$, for given $\mathbf{y} \in \mathbb{F}_q^b$, is invariant under scalar multiplication the rank of the matrix $B(\tilde{\mathbf{y}})$ with $\tilde{\mathbf{y}} = (\tilde{y_1} : ... : \tilde{y_b}) \in \mathbb{P}^{b-1}(\mathbb{F}_q)$ is well defined. Under the Lazard correspondence we may identify $1 \neq g \in G'$ with an element $\mathbf{g} \in \mathfrak{g}' \cong \mathbb{F}_q^b$, the associated Lie algebra, and similarly identify $\mathbf{g}$ with $\tilde{\mathbf{g}} \in \mathbb{P}^{b-1}(\mathbb{F}_q)$. Since the matrix B is skew-symmetric its determinant $\det(B)$ is a square whose square root $\mathrm{Pf}(B) := \sqrt{\det(B)}$ is the *Pfaffian* of B. If a is odd, then $\mathrm{Pf}(B) = 0$. Assume that $\mathrm{Pf}(B) \neq 0$. Then $\mathrm{Pf}(B)$ defines a hypersurface in $\mathbb{P}^{b-1}$ and the $\mathbb{F}_q$-rational points $\tilde{\mathbf{y}}$ of this hypersurface correspond to matrices $B(\tilde{\mathbf{y}})$ of a certain non-maximal rank. The $\mathbb{F}_q$-rational points which do not lie on the hypersurface will be of maximal rank a. We refer to points $\tilde{\mathbf{y}} \in \mathbb{P}^{b-1}(\mathbb{F}_q)$ as being 'of rank 2i' for some i if the associated matrix $B(\tilde{\mathbf{y}})$ is of rank $2i$. The condition $g \in \mathrm{Ker}(\omega_{\mathbf{y}})$ in the definition of $K_G^i(g)$ (see equation (5)) defines a hyperplane H_g given by the dot product $\tilde{\mathbf{g}}.\tilde{\mathbf{y}} = 0$ in $\mathbb{P}^{b-1}(\mathbb{F}_q)$. We may thus talk about the 'hyperplane defined by g' as H_g. The numbers $K_G^i(g)$ are simply the number of $\mathbb{F}_q$-rational points of this hyperplane which intersect the hypersurface defined by $\mathrm{Pf}(B)$ in $\mathbb{P}^{b-1}(\mathbb{F}_q)$ giving rise to a point 'of rank $2i$'.

We will now proceed with some examples. In each case it turns out that the matrices $B(\mathbf{Y})$ have the form

$$B(\mathbf{Y}) = \begin{pmatrix} 0 & U(\mathbf{Y}) \\ -U(\mathbf{Y})^{\mathrm{tr}} & 0 \end{pmatrix}$$

where $U(\mathbf{Y})$ is a matrix.

Example 4.1 (Heisenberg group, $\mathrm{H}(\mathbb{F}_q)$)**.** Suppose that G is the Heisenberg group $\mathrm{H}(\mathbb{F}_q)$. The matrix $U(\mathbf{Y})$, where $\mathbf{Y} = (Y_1)$ has a single variable, is simply the 1×1 matrix with entry Y_1. There are two cases, $i = 0$ and $i = 1$, for the rank of the matrix $U(\mathbf{Y})$. The number of vectors $\mathbf{y}$ in each case is 1 and $q - 1$ respectively. When $g = 1$ is the identity, we have $K_G^0(1) = 1$ and $V_G^0(1) = 0$ so that $\zeta_G^0(s, 1) = q^2$. We also have $K_G^1(1) = q - 1$ and $V_G^1(1) = 0$ so that $\zeta_G^1(s, 1) = q^{1-s}(q - 1)$. Together, this gives us

$$\zeta_G(s, 1) = q^2 + q^{-s+1}(q - 1)$$

and when $s = 1$ this is simply $k(G)$, the class number.

Suppose now that $1 \neq g \in G'$. This occurs with multiplicity $(q-1)$. It is not hard to see that $\zeta_G^0(s,g) = q^2$ since $K_G^0(g) = 1$ and $V_G^0(g) = 0$. Also, $K_G^1(g) = 0$ and $V_G^1(g) = q-1$ so that

$$\zeta_G(s,g) = q^2 - q^{-s+1}.$$

Example 4.2 (A quadric surface in $\mathbb{P}^2(\mathbb{F}_q)$)**.** Let $\mathfrak{g}$ be the 7-dimensional nilpotent $\mathbb{F}_q$-Lie algebra of class 2 with $\mathbb{F}_q$-basis $(x_1, ..., x_4, y_1, y_2, y_3)$ subject to the relations $[x_1, x_3] = y_1$, $[x_1, x_4] = y_2$, $[x_2, x_3] = y_3$, $[x_2, x_4] = y_1$. With respect to this basis we have

$$U(\mathbf{Y}) = \begin{pmatrix} Y_1 & Y_2 \\ Y_3 & Y_1 \end{pmatrix}.$$

The determinant of this matrix is defines the quadric surface $Y_1^2 - Y_2Y_3$ in $\mathbb{P}^2(\mathbb{F}_q)$. There are three cases, $i = 0, 1$ and 2, for the rank of the matrix $U(\mathbf{Y})$. The number of such vectors $\mathbf{y}$ in each case is 1, $(q+1)(q-1)$ and $q^2(q-1)$ respectively. As usual, the $i = 0$ case corresponds to when $\mathbf{y}$ is zero, $i = 1$ when $\tilde{\mathbf{y}}$ lies on the curve and $i = 2$ otherwise. When g is the identity we can compute

$$K_G(1) = (1, (q+1)(q-1), q^2(q-1))$$

and $V_G(1) = (0, 0, 0)$ so that $k(G) = q^4 + q^2(q+1)(q-1) + q^2(q-1)$.

Now suppose that $g \neq 1$. We have two cases. The first case is when H_g corresponds to a tangent of the surface $Y_1^2 = Y_2Y_3$. This occurs with multiplicity $(q+1)(q-1)$. In this case

$$K_G(g) = (1, (q-1), q(q-1))$$

and

$$V_G(g) = (0, q(q-1), (q^2-q)(q-1)).$$

The second case is when H_g corresponds to a line which intersects the surface $Y_1^2 = Y_2Y_3$ at two distinct points. This occurs with multiplicity $q^2(q-1)$. In this case

$$K_G(g) = (1, 2(q-1), (q-1)(q-1))$$

and

$$V_G(g) = (0, (q-1)(q-1), (q^2-q+1)(q-1)).$$

Example 4.3 (A quadric surface in $\mathbb{P}^3(\mathbb{F}_q)$)**.** Let $\mathfrak{g}$ be the 8-dimensional nilpotent $\mathbb{F}_q$-Lie algebra of class 2 with $\mathbb{F}_q$-basis $(x_1, ..., x_4, y_1, ..., y_4)$ subject to the relations $[x_1, x_3] = y_1$, $[x_1, x_4] = y_2$, $[x_2, x_3] = y_3$, $[x_2, x_4] = y_4$. With respect to this basis we have

$$U(\mathbf{Y}) = \begin{pmatrix} Y_1 & Y_2 \\ Y_3 & Y_4 \end{pmatrix}.$$

The determinant of this matrix defines the quadric surface $Y_1Y_4 - Y_2Y_3$ in $\mathbb{P}^3(\mathbb{F}_q)$. There are three cases, $i = 0, 1$ and 2, for the rank of the matrix $U(\mathbf{Y})$. The number of vectors $\mathbf{y}$ in each case is 1, $(q+1)^2(q-1)$ and $(q^3-q)(q-1)$ respectively. As usual, the $i = 0$ case corresponds to when $\mathbf{y}$ is zero, $i = 1$ when $\tilde{\mathbf{y}}$ lies on the curve and $i = 2$ otherwise. When g is the identity we compute

$$K_G(1) = (1, (q+1)^2(q-1), q^4 - 1 - (q+1)^2(q-1))$$

and $V_G(1) = (0, 0, 0)$ so that $k(G) = q^4 + q^2(q+1)^2(q-1) + q^4 - 1 - (q+1)^2(q-1)$.

Now suppose that $g \neq 1$. We have two cases. The first case is when g corresponds to a tangent of the surface $Y_1Y_4 = Y_2Y_3$. This occurs with multiplicity $(q+1)^2(q-1)$. In this case

$$K_G(g) = (1, (2q+1)(q-1), (q^2-q)(q-1))$$

and

$$V_G(g) = (0, q^2(q-1), (q^3-q^2)(q-1)).$$

The second case is when g corresponds to a plane which intersects the surface $Y_1Y_4 = Y_2Y_3$ and is non-tangent. This occurs with multiplicity $(q^3-q)(q-1)$. In this case

$$K_G(g) = (1, (q+1)(q-1), q^2(q-1))$$

and

$$V_G(g) = (0, q(q+1)(q-1), (q^3-q^2-q)(q-1)).$$

The following examples were studied by Boston & Isaacs [2].

Example 4.4 (Elliptic curves in $\mathbb{P}^2(\mathbb{F}_p)$)**.** Let p be a prime and $\alpha \in \mathbb{F}_p^*$. Let $\mathfrak{g}_\alpha$ be the 9-dimensional nilpotent $\mathbb{F}_p$-Lie algebra of class 2 with $\mathbb{F}_p$-basis $(x_1, ..., x_6, y_1, y_2, y_3)$ subject to the relations $[x_1, x_4] = y_1$, $[x_1, x_5] = y_2$, $[x_1, x_6] = \alpha y_3$, $[x_2, x_4] = y_3$, $[x_2, x_5] = y_1$, $[x_2, x_6] = y_2$, $[x_3, x_4] = y_3$, $[x_3, x_6] = y_1$. With respect to this basis we have

$$U(\mathbf{Y}) = \begin{pmatrix} Y_1 & Y_2 & \alpha Y_3 \\ Y_3 & Y_1 & Y_2 \\ Y_3 & 0 & Y_1 \end{pmatrix}.$$

The determinant of this matrix defines an elliptic curve E_α in $\mathbb{P}^2(\mathbb{F}_p)$ and let n_α denote the number of $\mathbb{F}_p$-rational points of the curve E_α. There are three cases, $i = 0, 1$ and 2, for the rank of the matrix $U(\mathbf{Y})$. The number of such vectors $\mathbf{y}$ in each case is 1, $n_\alpha(p-1)$ and $(p^2+p+1-n_\alpha)(p-1)$ respectively. As usual, the $i = 0$ case corresponds to when $\mathbf{y}$ is zero, $i = 1$ when $\tilde{\mathbf{y}}$ lies on the curve and $i = 2$ otherwise. When g is the identity we can compute

$$K_G(1) = (1, n_\alpha(p-1), (p^2+p+1-n_\alpha)(p-1))$$

and $V_G(1) = (0, 0, 0)$ so that $k(G) = p^6 + p^2 n_\alpha(p-1) + (p^2+p+1-n_\alpha)(p-1)$.

Now suppose that $g \neq 1$. Let k_α denote the number of inflection points of E_α. We have four cases depending on how many times the line H_g corresponding to g intersects the elliptic curve E_α at rational points, this can be zero, once, twice or three times occurring with multiplicities $p^2+p+1-n_\alpha(p+1)+\frac{2}{3}\binom{n_\alpha}{2}+\frac{n_\alpha-k}{3}$, $n_\alpha(p+1)-(n_\alpha-k)-\binom{n_\alpha}{2}$, $(n_\alpha-k)(p-1)$ and $\binom{n_\alpha}{2}$ respectively.

Suppose that $1 \neq g$ defines a line which intersects the curve at m rational points where $m = 0, 1, 2, 3$. We have

$$K_G(g) = (1, m(p-1), (p+1-m)(p-1))$$

and

$$V_G(g) = (0, (n_\alpha-m)(p-1), (p^2+m-n_\alpha)(p-1)).$$

5 Acknowledgements

I would like to thank Christopher Voll for his support, guidance and insight throughout this project and for numerous helpful discussions.

References

[1] M. Abért. On the probability of satisfying a word in a group. *J. Group Theory*, 9(5):685–694, 2006.

[2] N. Boston and I. M. Isaacs. Class numbers of p-groups of a given order. *J. Algebra*, 279(2):810–819, 2004.

[3] M. Boyarchenko and M. Sabitova. The orbit method for profinite groups and a p-adic analogue of brown's theorem. *Israel J. Math*, 165:67–91, 2008.

[4] J. D. Dixon, L. Pyber, Á. Seress, and A. Shalev. Residual properties of free groups and probabilistic methods. *J. reine angew. Math. (Crelle's)*, 556:159–172, 2003.

[5] S. Garion and A. Shalev. Commutator maps, measure preservation and T-systems. *Trans. Amer. Math. Soc.*, 361(9):4631–4351, 2009.

[6] J. González-Sánchez. Kirillov's orbit method for p-groups and pro-p groups. *Comm. Algebra*, 37(12):4476–4488, 2009.

[7] I. M. Isaacs. *Character Theory of Finite Groups*, volume 359. American Math. Soc., 1976.

[8] A. Jaikin-Zapirain. Zeta function of representations of compact p-adic analytic groups. *J. Amer. Math. Soc.*, (19):91–118, 2006.

[9] E. I. Khukrho. *p-automorphisms of finite p-groups*, volume 246 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1198.

[10] M. Larsen and A. Shalev. Fibres of word maps and some applications. *J. Algebra*, 354:36–48, 2012.

[11] M. Levy. On the probability of solving a word in nilpotent groups of class 2. (arχiv 1101.4286v1).

[12] E. A. O'Brien and C. Voll. Enumerating classes and characters of p-groups. *Trans. Amer. Math. Soc.*, 367:7775–7796, 2015.